

12

EUROPEAN PATENT APPLICATION

21 Application number: 87810687.1

51 Int. Cl.⁴: **C 07 H 19/173**
A 61 K 31/70

22 Date of filing: 23.11.87

30 Priority: 27.11.86 JP 283904/86

43 Date of publication of application:
01.06.88 Bulletin 88/22

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **NIPPON ZOKI PHARMACEUTICAL CO. LTD.**
10, 2-chome Hirano-machi Higashi-ku
Osaka-shi Osaka-fu (JP)


72 Inventor: **Yamada, Toshio c/o Institute of Bio-Active Science**
Nippon Zoki Pharm. Co. Ltd. 442-1, Kinashi
Yashiro-cho Kato-gun Hyogo (JP)

Kageyama, Ken-ichi Institute of Bio-Active Science
Nippon Zoki Pharm. Co. Ltd. 442-1, Kinashi
Yashiro-cho Kato-gun Hyogo (JP)

74 Representative: **Kerr, Andrew**
Postfach 122 Finkelerweg 44
CH-4144 Arlesheim BL (CH)

54 Novel adenosine derivatives and pharmaceutical compositions containing them as an active ingredient.

57 The present invention relates to novel adenosine derivatives having the formula (I):



wherein each of R₁, R₂ and R₃, which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl group;

and pharmaceutically acceptable salt thereof. These compounds are useful as antihypertensive agents, for example, for the treatments of hypertension and various diseases caused by hypertension, such as cerebrovascular diseases, cardiopathy or renal insufficiency.

Description

NOVEL ADENOSINE DERIVATIVES AND PHARMACEUTICAL COMPOSITION CONTAINING THEM AS AN ACTIVE INGREDIENT


5 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel adenosine derivatives, pharmaceutically acceptable salts thereof and pharmaceutical compositions containing them as an active ingredient.

Hypertension is one of the most serious risk factors causing cerebral apoplexy and cardiopathy which are included in the major causes of death. The frequency of hypertension is increasing as becoming old, and it is
10 estimated that about 20% persons of the total population in Japan are suffering from hypertension. Thus various antihypertensive agents such as diuretics, sympathetic depressant drugs, direct-acting vasodilators, calcium antagonists or angiotensin antagonists have been developed and used for a treatment of hypertension. However, for example, side effects such as bradycardia causing various symptoms such as tiredness, depression of activity, disturbance of cerebral circulation or cerebral ischemia are sometimes
15 observed under the treatment with sympathetic depressant drugs. Therefore, the developments of antihypertensive agents having greater safety and effectiveness are desired. As a result of investigations for orally administrable antihypertensive compounds, the inventors have found adenosine derivatives having vasodilatively hypotensive effect as well as low toxicity and good safety.

An object of the present invention is to provide new adenosine derivatives and pharmaceutically acceptable salts thereof having an excellent antihypertensive effect. Another object of the present invention is to provide a process for preparing these adenosine derivatives. A further object of the invention is to provide pharmaceutical compositions containing these adenosine derivatives or pharmaceutically acceptable salts thereof as an active ingredient.

The adenosine derivatives of the present invention are represented by the following general formula (I):



wherein each of R_1 , R_2 and R_3 , which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl group.

Each of R_1 , R_2 and R_3 , which may be the same or different, represents hydrogen or a lower alkyl group, preferably a straight or branched alkyl group having 1 to 5 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl or tert-pentyl.

X represents hydrogen; a lower alkyl group, preferably a straight or branched alkyl group having 1 to 5 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl or tert-pentyl; an amino group; or halogen such as fluoride, chloride, bromide or iodide.

Y represents hydrogen or a lower alkyl group, preferably a straight or branched alkyl group having 1 to 5 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl or tert-pentyl.

The adenosine derivatives of the present invention include pharmaceutically acceptable salts of the compounds having formula (I) above, for example, salts with alkali metal such as sodium or potassium, with alkaline-earth metal such as calcium or barium, or with other metals such as aluminum, or salts with an organic base such as ammonia or the like. These salts can be prepared from free adenosine derivatives or other salts of these derivatives by a known method.

When optical isomers exist in the compounds of the invention, the present invention includes any of the dl, d

and l-isomers.

The adenosine derivatives of the present invention can be prepared as follows.

(1) Adenosine or adenosine derivatives having a lower alkyl group, an amino group or halogen at the 2'-position were alkylated at the 2'-O- or 3'-O-position by an alkylating agent to give the compounds of the present invention. A diazoparaffin, such as diazomethane, diazoethane, diazopropane or diazobutane, can be used as the alkylating agent. The appropriate solvent which does not inhibit the reaction such as 1,2-dimethoxyethane can be preferably used. This O-alkylating reaction can be carried out as follows: (i) The reaction mixture is reacted for several minutes to several hours at room temperature in the presence of a catalyst such as p-toluenesulfonic acid. (ii) The starting material is dissolved in about 80°C hot water and the alkylating agent such as diazoparaffin is added thereto, and the reaction mixture is reacted for several hours to a day.

(2) Both 3'-O- and 5'-O-positions of the adenosine derivatives are protected by tetraisopropylidisiloxane (TIPDS) group to carry out O-alkylation selectively at the 2'-O-position. A 6-chloropurine-9-riboside and TIPDS•Cl₂ (1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane dichloride) are stirred for several hours at room temperature to protect the 3'-O- and 5'-O-positions, and then 2'-O-position of the compound protected by TIPDS can be selectively alkylated by an alkylating agent such as methyl iodide, ethyl iodide, propyl iodide or butyl iodide in the presence of catalyst such as silver oxide. After the 2'-O-alkylation, an amination or alkylation at the 6-position can be carried out by reacting with ammonia or alkylamine such as methylamine, ethylamine, propylamine or butylamine with heating. The protecting group, TIPDS, can be removed by a conventional method to give the compounds of the present invention.

(3) In the similar manner, both 2'-O- and 3'-O-positions of the adenosine derivatives are protected by isopropylidene group to carry out O-alkylation selectively at the 5'-O-position. Namely, A 6-chloropurine-9-riboside and 2,2-dimethoxypropane are reacted for several hours at room temperature in the presence of catalyst such as p-toluenesulfonic acid to carry out isopropylidenation. After the 5'-O-alkylation, an amination or alkylation at 6-position can be carried out as mentioned above. The protecting group, isopropylidene group, can be removed by a conventional method, for example, treatment with formic acid, to give the compounds of the present invention.

The resulting compounds of the present invention can be purified by known methods such as distillation chromatography and recrystallization. Identification is established through, inter alia, melting point, elemental analysis, IR, NMR, UV, mass spectrum, etc.

Example

The following examples, which are illustrative only and not intended to limit the scope of the invention, describe the preparation of the compounds of the present invention.

Example 1.

[METHOD 1]

10 g of adenosine was suspended in 2 l of 1 mmole methanol solution of tin chloride dihydrate. 500 ml of 0.4-0.5M 1,2-dimethoxyethane solution of diazomethane was added to the solution with stirring. After stirring for 2 hours at room temperature, the solution was evaporated to dryness under reduced pressure. The residue was purified by ion exchange column chromatography and recrystallized from ethanol to give 2'-O-methyladenosine (Compound 1) and 3'-O-methyladenosine (Compound 2).

-Compound 1-

yield: 33.5 %

m.p.: 205.5 - 206 °C

NMR(D₂O): δ=3.43(3H,s), 3.93(1H,dd,J=2.4, 12.7Hz), 3.86(1H,dd,J=2.9, 12.7Hz), 4.26(1H,m), 4.40(1H,dd,J=5.9, 5.9Hz), 4.56(1H,dd,J=4.9, 5.9Hz), 6.01(1H,d,J=5.9Hz), 8.01(1H,s), 8.20(1H,s)

-Compound 2-

yield: 43.4 %

m.p.: 177.5 - 178 °C

NMR(D₂O): δ=3.54(3H,s), 3.83(1H,dd,J=2.9, 13.2Hz), 3.94(1H,dd,J=3.4, 13.2Hz), 4.11(1H,dd,J=4.4, 4.4Hz), 4.37(1H,m), 4.87(1H,dd,J=5.9, 5.9Hz), 6.00(1H,d,J=5.9Hz), 8.11(1H,s), 8.25(1H,s)

[METHOD 2]

5 g of adenosine was dissolved in 160 ml of 80°C hot water, and 500 ml of 1,2-dimethoxyethane solution of diazomethane was added thereto with stirring. The solution was cooled to room temperature with stirring and was evaporated to dryness under reduced pressure. The starting material, adenosine, was removed by ODS column chromatography to give a mixture of Compound 1 and Compound 2. 2.6 g of the mixture was separated by ion exchange column chromatography to obtain Compound 1 and Compound 2.

Example 2.

(i) 10 g of inosine was treated in the same manner as Example 13 (i) and (ii) mentioned later to give 12 g of 6-chloro-9-(2,3,5-O-triacetyl-β-D-ribofuranosyl)-9H-purine (yield: 78%).

(ii) 200 ml of 20% ammonia methanol solution was added to the resulting product, and the solution was stirred for 4 hours at room temperature. The solution was concentrated under reduced pressure and ethyl acetate was added thereto to precipitate 5.6 g of 6-chloro-9-β-D-ribofuranosyl-9H-purine (yield: 69%).

(iii) 4 g of 6-chloro-9-β-D-ribofuranosyl-9H-purine was suspended in 70 ml of pyridine. After 4.5 g of TIPDS·Cl₂ was added thereto and the solution was stirred for 2 hours at room temperature, the reaction mixture was purified by silica gel column chromatography to give 6.8 g of 6-chloro-9-(3,5-O-TIPDS-β-D-ribofuranosyl)-9H-purine (yield: 92%).

(iv) 2 g of the resulting product was dissolved in 70 ml of benzene, and 70 ml of ethyl iodide and 3 g of silver oxide were added to the solution. The solution was refluxed with heating. The reactant was checked every thirty minutes and ethyl iodide and silver oxide were added thereto. After filtering off silver iodide, the filtrate was concentrated to dryness, and the residue was dissolved in 20% ammonia methanol solution. The solution was heated for 6 hours at 80°C in an autoclave and purified by silica gel column chromatography to give 1.8 g of 2'-O-ethyl-3'-5'-O-TIPDS-adenosine.

(v) 1.5 g of the resulting product, 1 g of potassium fluoride and 3.7 g of tetraethylammonium chloride were dissolved in 65 ml of acetonitrile, and the solution was stirred for 15 hours at room temperature. After desalting, the solution was purified by ion exchange column chromatography to give 575 mg of 2'-O-ethyladenosine (Compound 3).

yield: 68 %

NMR(D₂O): δ = 1.09(3H,dd,J=6.8, 6.8Hz), 3.52-3.72(2H,m), 3.84(1H,dd,J=3.4, 12.7Hz), 3.91(1H,dd,J=2.9, 12.7Hz), 4.31(1H,m), 4.56(1H,dd,J=3.4, 5.4Hz), 4.64(1H,dd,J=5.4, 6.4Hz), 6.11(1H,d,J=6.4Hz), 8.25(1H,s), 8.34(1H,s)

Using butyl iodide instead of ethyl iodide, 2'-O-n-butylation was carried out in the same manner to give 2'-O-butyladenosine (Compound 4).

NMR(D₂O): δ = 0.64(3H,t,J=7.3Hz), 1.05(2H,m), 1.34(2H,m), 3.47(1H,m), 3.64(1H,m), 3.84(1H,dd,J=3.4, 12.7Hz), 3.92(1H,dd,J=2.9, 12.5Hz), 4.32(1H,m), 4.53(1H,dd,J=2.4, 5.4Hz), 4.62(1H,dd,J=5.4, 6.8Hz), 6.08(1H,d,J=6.84Hz), 8.24(1H,s), 8.34(1H,s)

Example 3.

[METHOD 1]

(i) 6 g of 5-amino-1-β-D-ribofuranosyl-4-imidazolecarboxamide was dissolved in 250 ml of 1N sodium ethoxide. 20 ml of ethyl acetate was added thereto, and the reaction mixture was heated for 3 hours at 120°C. After the solution was neutralized and desalted, 70 ml of acetic anhydride and 70 ml of pyridine were added thereto and the solution was stirred for 12 hours at room temperature. The solvent was distilled off under reduced pressure to give 9 g of 2-methyl-2',3',5'-O-triacetylinosine (yield: 95%).

(ii) 10.3 g of the resulting product was dissolved in 140 ml of chloroform. 14 ml of thionyl chloride and 3.5 ml of dimethylformamide were added thereto, and the solution was refluxed with heating for 2 hours. The reaction mixture was poured into ice cold water, and the separated chloroform layer was washed to be neutralized. After drying over sodium sulfate anhydride, the solvent was distilled off to give 9.5 g of 6-chloro-2-methyl-9-(2,3,5-O-triacetyl-β-D-ribofuranosyl)-9H-purine (yield: 88.7%).

(iii) The resulting product was dissolved in 150 ml of 20% ammonia methanol solution and heated for 8 hours in an autoclave. The solution was concentrated to dryness under reduced pressure and desalted to give 5.5 g of 2-methyladenosine (yield: 87.9%).

(iv) 2-methyladenosine was methylated in the same manner as Example [METHOD 1] to give 1.8 g of 2,2'-O-dimethyladenosine (Compound 5) and 2.9 g of 2,3'-O-dimethyladenosine (Compound 6).

-Compound 5-

yield: 34.3 %

m.p.: 169 - 170 °C

NMR(D₂O): δ = 2.47(3H,s), 3.39(3H,s), 3.83(1H,dd,J=2.44, 12.7Hz), 3.91(1H,dd,J=2.44, 12.7Hz), 4.32(1H,m), 4.52(1H,dd,J=4.9, 6.8Hz), 4.60(1H,dd,J=2.0, 4.9Hz), 6.05(1H,d,J=6.8Hz), 8.21(1H,s)

Compound 6-

yield: 55.2 %

m.p.: 203 - 203.5 °C

NMR(D₂O): δ = 2.50(3H,s), 3.55(3H,s), 3.84(1H,dd,J=2.9, 12.7Hz), 3.96(1H,dd,J=2.4, 12.7Hz), 4.13(1H,dd,J=2.9, 5.4Hz), 4.42(1H,m), 4.91(1H,dd,J=5.4, 6.4Hz), 6.00(1H,d,J=6.4), 8.23(1H,s)

[METHOD 2]

10 g of 5-amino-1-β-D-ribofuranosyl-4-imidazolecarbonitrile was dissolved in 20% ammonia methanol solution. 10 ml of acetonitrile was added thereto and the solution was heated for 5 hours at 180°C. The solvent was distilled off, purified by silica gel column chromatography and recrystallized from methanol to give 4.2 g of

2-methyladenosine (yield: 35.9%). 2-methyladenosine was methylated in the same manner as the above-mentioned [METHOD 1] to give Compound 5 and Compound 6.

Example 4.

Using isobutylnitrile instead of acetonitrile, 2-isopropyl-2'-O-methyladenosine (Compound 7) and 2-isopropyl-3'-O-methyladenosine (Compound 8) were obtained in the same manner as Example 2 [METHOD 2].

-Compound 7-

NMR(D₂O): δ = 1.25(6H,d,J=7Hz), 2.99(1H,sep.,J=7Hz), 3.39(3H,s), 3.83(1H,dd,J=3.3, 12.9Hz), 3.91(1H,dd,J=2.6, 12.9Hz), 4.29(1H,m), 4.58(1H,dd,J=4.9, 6.4Hz), 4.63(1H,dd,J=2.4, 4.9Hz), 6.07(1H,d,J=6.4Hz)

-Compound 8-

NMR(D₂O): δ = 1.35(6H,d,J=6.9Hz), 3.14(1H,sep.,J=6.9Hz), 3.54(3H,s), 3.84(1H,dd,J=4.1, 12.8Hz), 3.94(1H,dd,J=3.0, 12.8Hz), 4.24(1H,dd,J=5.4, 5.4Hz), 4.36(1H,m), 5.03(1H,dd,J=5.4, 5.4Hz), 6.08(1H,d,J=5.4Hz), 8.34(1H,s)

Example 5.

6-chloro-2-methyl-9- β -D-ribofuranosyl-9H-purine was treated in the same manner as Example 2 to give 6-chloro-2-methyl-9-(3,5-O-TIPDS- β -D-ribofuranosyl)-9H-purine. After the 2'-position of the compound was O-ethylated by using ethyl iodide, the 6-position of the compound was aminated by using ammonia. The protecting group, TIPDS, was removed to give 2-methyl 2'-O-ethyladenosine (Compound 9).

NMR(D₂O): δ = 1.07(3H,dd,J=6.8, 6.8Hz), 2.52(3H,s), 3.38(3H,s), 3.48-3.57(1H,m), 3.64-3.73(1H,m), 3.86(1H,dd,J=2.9, 12.7Hz), 3.94(1H,dd,J=2.4, 12.7Hz), 4.35(1H,ddd,J=2.0, 2.4, 2.9Hz), 4.59(1H,dd,J=2.0, 4.9Hz), 4.68(1H,dd,J=4.9, 7.3Hz), 6.07(1H,d,J=7.3Hz), 8.25(1H,s)

Using butyl iodide instead of ethyl iodide, 2'-O-butylation was carried out to give 2-methyl-2'-O-n-butyladenosine (Compound 10) in the same manner.

NMR(CDCl₃): δ = 0.79(3H,t,J=7.3Hz), 1.22(2H,qt,J=7.3, 7.3Hz), 1.39(2H,ddt,J=7.3, 7.3, 7.3Hz), 2.56(3H,s), 3.33(1H,m), 3.47(1H,m), 3.74(1H,ddd,J=1.5, 13.7, 14.2Hz), 3.96(1H,ddd,J=1.5, 2.0, 13.7Hz), 4.35(1H,m), 4.51(1H,d,J=4.4Hz), 4.78(1H,dd,J=4.4, 7.8Hz), 5.64(2H,brs,D₂O exchangeable), 5.78(1H,d,J=7.8Hz), 7.02(1H,dd,J=2.0, 14.2Hz,D₂O-exchangeable), 7.74(1H,s)

Example 6.

14 ml of 2,2-dimethoxypropane, 16 ml of acetone and 1.3 g of p-toluenesulfonic acid were added to 1.5 g of 6-chloro-9- β -D-ribofuranosyl-9H-purine, and the reaction mixture was stirred for 2 hours at room temperature to give 6-chloro-9-(2,3-O-isopropylidene- β -D-ribofuranosyl)-9H-purine. The solution was concentrated to dryness and separated with chloroform/water. The chloroform layer was washed and dried over sodium sulfate anhydride. After the solvent was distilled off, the residue was dissolved in 30 ml of acetone. 30 ml of methyl iodide and 1.2 g of silver oxide were added to the solution and 5'-O-methyl-2',3'-O-isopropylideneadenosine was obtained in the same manner as Example 2 (iv). After filtration, the solution was concentrated to dryness, and the residue was dissolved in 20% ammonia methanol solution to be aminated. The solvent was distilled off and 90% formic acid was added thereto, and then the solution was stirred for 30 minutes at room temperature to remove the protecting group. The solution was concentrated to dryness under reduced pressure and recrystallized from ethanol to give 1.1 g of 5'-O-methyladenosine (Compound 11).

NMR(D₂O): δ = 3.43(3H,s), 3.71-3.82(2H,m), 4.33(1H,m), 4.43(1H,dd,J=5.4, 5.4Hz), 4.78(1H,dd,J=5.4, 5.4Hz), 6.12(1H,d,J=5.4Hz), 8.31(1H,s), 8.41(1H,s)

Using butyl iodide instead of methyl iodide, 5'-O-butylation was carried out to give 5'-O-n-butyladenosine (Compound 12) in the same manner.

NMR(D₂O): δ = 0.94(3H,t,J=7.3Hz), 1.41(2H,qt,J=7.3, 7.3Hz), 1.60(2H,tt,J=7.3, 7.3Hz), 3.54(2H,m), 3.66(1H,dd,J=3.4, 11.2Hz), 3.78(1H,dd,J=2.9, 11.2Hz), 4.19(1H,m), 4.34(1H,dd,J=4.9, 4.9Hz), 4.55(1H,dd,J=4.4, 4.9Hz), 6.05(1H,d,J=4.4Hz), 8.19(1H,s), 8.40(1H,s)

Example 7.

In the same manner as Example 1 [METHOD 1], 5'-O-methyladenosine was methylated by using diazomethane and separated by ion exchange column chromatography to give 2',5'-O-dimethyladenosine (Compound 13) and 3',5'-O-dimethyladenosine (Compound 14).

-Compound 13-

NMR(D₂O): δ = 3.41(3H,s), 3.46(3H,s), 3.67-3.76(2H,m), 4.27(1H,m), 4.40(1H,dd,J=4.4, 4.9Hz), 4.52(1H,dd,J=4.9, 5.4Hz), 6.07(1H,d,J=5.4Hz), 6.07(1H,d,J=5.4Hz), 8.10(1H,s), 8.26(1H,s)

-Compound 14-

NMR(D₂O): δ = 3.42(3H,s), 3.47(3H,s), 3.68-3.77(2H,m), 4.30(1H,m), 4.50(1H,dd,J=4.9, 5.4Hz), 4.57(1H,dd,J=4.4, 4.9Hz), 6.2(1H,d,J=5.4Hz), 8.23(1H,s), 8.35(1H,s)

Example 8.

In the same manner as Example 6, 6-chloro-9- β -D-ribofuranosyl-9H-purine was isopropylidenated, and methylated at the 5'-O-position by using methyl iodide. After the 6-position of the compound was aminated by ammonia, the protecting group was removed to give 2,5'-O-dimethyladenosine (Compound 15).

NMR(D₂O): δ =2.55(3H,s), 3.41(3H,s), 3.73-3.78(2H,m), 4.32(1H,m), 4.42(1H,dd,J=4.9, 4.9Hz), 4.77(1H,dd,J=4.9, 4.9Hz), 6.07(1H,d,J=4.9Hz), 8.25(1H,s)

Using butyl iodide instead of methyl iodide, 5'-O-butylation was carried out to give 2-methyl-5'-O-n-butyladenosine (Compound 16) in the same manner.

NMR(CD₃OD): δ =0.95(3H,t,J=7.3Hz), 1.40(2H,qt,J=7.3, 7.3Hz), 1.60(2H,tt,J=7.3, 7.3Hz), 2.51(3H,s), 3.50-3.70(3H,m), 3.78(1H,dd,J=2.9, 10.7Hz), 4.17(1H,m), 4.34(1H,dd,J=4.9, 4.9Hz), 4.52(1H,dd,J=4.4, 4.9Hz), 6.00(1H,d,J=4.4Hz), 8.11(1H,s)

Example 9.

6-chloro-9- β -D-ribofuranosyl-9H-purine was treated in the same manner as Example 2 to give 6-chloro-9-(3,5-O-TIPDS- β -D-ribofuranosyl)-9H-purine. After the compound was 2'-O-methylated, the 6-position of the compound was methylaminated by methylamine to give N⁶-2'-O-dimethyladenosine (Compound 17).

NMR(D₂O): δ =3.00(3H,s), 3.40(3H,s), 3.81(1H,dd,J=3.4, 12.7Hz), 3.89(1H,dd,J=2.9, 12.7Hz), 4.26(1H,ddd,J=2.9, 3.4, 3.4Hz), 4.44(1H,dd,J=5.4, 6.4Hz), 4.57(1H,dd,J=3.4, 5.4Hz), 6.03(1H,d,J=6.4Hz), 8.10(1H,s), 8.19(1H,s)

Using butylamine instead of methylamine, N⁶-n-butyl-2'-O-methyladenosine (Compound 18) was obtained in the same manner.

NMR(CDCl₃): δ =0.95(3H,t,J=7.3Hz), 1.43(2H,qt,J=7.3, 7.3Hz), 1.65(2H,tt,J=7.3, 7.3Hz), 3.32(3H,s), 3.61(3H,m), 3.73(1H,m), 3.94(1H,m), 4.33(1H,m), 4.57(1H,d,J=4.4Hz), 4.73(1H,dd,J=4.4, 7.8Hz), 5.81(1H,d,J=7.8Hz), 5.93(1H,brs,D₂O-exchangeable), 6.96(1H,brs,D₂O-exchangeable), 7.74(1H,s), 8.32(1H,s)

Example 10.

In the same manner as Example 6, 6-chloro-9- β -D-ribofuranosyl-9H-purine was isopropylidenated. After 5'-O-methylation, the 6-position of the compound was methylaminated by methylamine to give N⁶-5'-O-dimethyladenosine (Compound 19).

NMR(D₂O): δ =3.05(3H,s), 3.40(3H,s), 3.68-3.79(2H,m), 4.29(1H,m), 4.38(1H,dd,J=4.9, 5.4Hz), 4.72(1H,dd,J=4.9, 5.4Hz), 6.03(1H,d,J=4.9Hz), 8.17(1H,s), 8.23(1H,s)

Using butylamine instead of methylamine, N⁶-n-butyl-5'-O-methyladenosine (Compound 20) was obtained in the same manner.

NMR(CDCl₃): δ =0.95(3H,t,J=7.3Hz), 1.44(2H,qt,J=7.3, 7.3Hz), 1.66(2H,tt,J=7.3, 7.3Hz), 3.33(3H,s), 3.54-3.68(4H,m), 4.35-4.45(3H,m), 5.87(1H,brs,D₂O-exchangeable), 5.93(1H,d,J=5.9Hz), 8.00(1H,s), 8.28(1H,s)

Example 11.

6-chloro-2-methyl-9- β -D-ribofuranosyl-9H-purine was treated in the same manner as mentioned above to give 6-chloro-2-methyl-9-(3,5-O-TIPDS- β -D-ribofuranosyl)-9H-purine or 6-chloro-2-methyl-9-(2,3-O-isopropylidene- β -D-ribofuranosyl)-9H-purine, and then the resulting compound was 2'-O-alkylated or 5'-O-alkylated. After the 6-position of the compound was aminated or alkylaminated by ammonia or alkylamine, the protecting group, TIPDS or isopropylidene, was removed to give the following compounds.

2,N⁶,2'-O-trimethyladenosine (Compound 21)

NMR(D₂O): δ =2.40(3H,s), 2.97(3H,s), 3.37(3H,s), 3.81(1H,dd,J=2.9, 12.7Hz), 3.90(1H,dd,J=2.9, 12.7Hz), 4.29(1H,m), 4.44(1H,dd,J=4.9, 6.8Hz), 4.57(1H,dd,J=2.4, 4.9Hz), 5.96(1H,d,J=6.8Hz), 8.06(1H,s)

2,N⁶-dimethyl-2'-O-ethyladenosine (Compound 22)

NMR(D₂O): δ =1.05(3H,dd,J=6.8, 6.8Hz), 2.50(3H,s), 3.09(3H,s), 3.37(3H,s), 3.46 3.74(2H,m), 3.85(1H,dd,J=2.9, 13.2Hz), 3.93(1H,dd,J=2.4, 13.2Hz), 4.34(1H,m), 4.56(1H,dd,J=1.5, 4.9Hz), 4.63(1H,dd,J=5.4, 6.8Hz), 6.02(1H,d,J=6.8Hz), 8.15(1H,s)

N⁶-n-butyl-2,2'-O-dimethyladenosine (Compound 23)

NMR(CDCl₃): δ =0.94(3H,t,J=7.3Hz), 1.43(2H,qt,J=7.3, 7.3Hz), 1.63(2H,tt,J=7.3, 7.3Hz), 2.54(3H,s), 3.32(3H,s), 3.62(2H,m), 3.74(1H,m), 3.95(1H,dd,J=1.5, 12.7Hz), 4.33(1H,m), 4.56(1H,d,J=4.4Hz), 4.74(1H,dd,J=4.4, 7.8Hz), 5.77(1H,d,J=7.8Hz), 5.79(1H,brs,D₂O-exchangeable), 7.33(1H,brs,D₂O-exchangeable), 7.66(1H,s)

2,N⁶,5'-O-trimethyladenosine (Compound 24)

NMR(D₂O): δ = 2.52(3H,s), 3.10(3H,s), 3.41(3H,s), 3.68-3.78(2H,m), 4.30(1H,m), 4.40(1H,dd,J = 4.9, 4.9Hz), 4.72(1H,dd,4.9, 4.9Hz), 6.05(1H,d,J = 4.9Hz), 8.20(1H,s)

N⁶-n-butyl-2,5'-O-dimethyladenosine (Compound 25)

NMR(CDCl₃): δ = 0.95(3H,t,J = 7.3Hz), 1.43(2H,qt,J = 7.3, 7.3Hz), 1.64(2H,tt,J = 7.3, 7.3Hz), 2.52(3H,s), 3.5-3.7(4H,m), 4.3-4.4(3H,m), 5.77(1H,brs,D₂O-exchangeable), 5.87(1H,d,J = 5.9Hz), 7.91(1H,s)

Example 12.

1 g of 2,2'-O-dimethyladenosine (Compound 5) was dissolved in 5 ml of dimethylformamide, and 2 ml of methyl iodide was added thereto and the solution was stirred for 30 to 40 minutes. After the solvent was distilled off, 5 ml of 0.5N sodium hydroxide was added and the solution was heated for 75 minutes at 100°C. The solution was neutralized and desalted by hydrophobic column chromatography, and then recrystallized from methanol to give 2,N⁶,2'-O-trimethyladenosine (Compound 21).

In the same manner, 2,3'-O-dimethyladenosine was used as a starting material to give 2,N⁶,3'-O-trimethyladenosine (Compound 26).

NMR(D₂O): δ = 2.46(3H,s), 3.05(3H,s), 3.56(3H,s), 3.84(1H,dd,J = 3.4, 12.7Hz), 3.96(1H,dd,J = 2.0, 12.7Hz), 4.11(1H,m), 4.41(1H,m), 4.86(1H,t-like, J = 5.9Hz), 5.94(1H,d,J = 5.9Hz), 8.10(1H,s)

Example 13

(i) 500 ml of acetic anhydride and 500 ml of pyridine were added to 45 g of guanosine, and the reaction mixture was stirred for 12 hours at room temperature. The precipitated crystalline was separated by filtration and washed with water. The filtrate was concentrated under reduced pressure, and the precipitated crystalline was obtained by filtration. The first and second crystals were put together to give 2',3',5'-O-triacetylguanosine (yield: 94.40%)

(ii) 375 ml of phosphorus oxychloride and 20 ml of diethylaniline was added to 55 g of 2',3',5'-O-triacetylguanosine, and the reaction mixture was refluxed with heating for 3 minutes. The solution was concentrated to 100 ml under reduced pressure, poured into ice-cold water and extracted with dichloromethane. The dichloromethane layer was washed to be neutralized and dried over sodium sulfate anhydride. After purification by silica gel column chromatography, the solution was concentrated under reduced pressure to give 40 g of 6-chloro-2-amino 9-(2,3,5-O-triacetyl- β -D-ribofuranosyl)-9H-purine (yield 69%).

(iii) 4.5 g of the resulting product was dissolved in 40 ml of cooled hydrochloric acid. 1.05 g of sodium nitrite dissolved in 3 ml of water was slowly added to the solution with stirring. The reaction mixture was neutralized by cooled ammonia water and extracted with dichloroethane. The dichloroethane layer was dried over sodium sulfate anhydride and concentrated to dryness to give 2,6-dichloro-9-(2,3,5-O-triacetyl- β -D-ribofuranosyl)-9H-purine.

(iv) The above residue was 100 ml of 20% ammonia methanol solution and heated for 8 hours at 70°C in an autoclave to give 2-chloroadenosine.

(v) In the same manner as Example 1 [METHOD 1], 2-chloroadenosine was methylated to give 2-chloro-2'-O-methyladenosine (Compound 27) and 2-chloro-3'-O-methyladenosine (Compound 28).

-Compound 27-

NMR(D₂O): δ = 3.43(3H,s), 3.84(1H,dd,J = 3.4, 13.2Hz), 3.92(1H,dd,J = 2.5, 13.2Hz), 4.30(1H,m), 4.40(1H,dd,J = 6.8, 4.9Hz), 4.60(1H,dd,J = 3.4, 4.9Hz), 6.05(1H,d,J = 6.8Hz), 8.28(1H,s)

-Compound 28-

NMR(D₂O): δ = 3.53(3H,s), 3.83(1H,dd,J = 3.4, 12.9Hz), 3.95(1H,dd,J = 2.6, 12.9Hz), 4.12(1H,dd,J = 5.9, 5.9Hz), 4.38(1H,m), 4.87(1H,dd,J = 5.9, 5.9Hz), 5.98(1H,d,J = 5.9Hz), 8.27(1H,s)

Example 14.

3 g of 2-amino-2'-O-methyladenosine, the Compound 30 in Example 15, was dissolved in 35 ml of 42% fluoboric acid. 1.3 g/10 ml aqueous solution of sodium nitrite was added to the solution with stirring at -5°C to -10°C and stirred for one hour. The reaction mixture was cooled to -20°C and neutralized by 50% sodium hydroxide aqueous solution. The solution was desalted by hydrophobic column chromatography and crystallized from methanol to give 697 mg of 2-fluoro-2'-O-methyladenosine (Compound 29).

Yield: 23 %

NMR(D₂O): δ = 3.45(3H,s), 3.84(1H,dd,J = 3.9, 13.2Hz), 3.91(1H,dd,J = 2.9, 13.2Hz), 4.27(1H,m), 4.50(1H,dd,J = 5.9, 6.4Hz), 4.60(1H,dd,J = 3.4, 5.9Hz), 6.06(1H,d,J = 6.4Hz), 8.29(1H,s)

Example 15.

The 6-position of 2-amino-6-chloro-9-(2,3,5-O-triacetyl- β -D-ribofuranosyl)-9H-purine was aminated by ammonia to give 2-aminoadenosine. In the same manner as Example 1 [METHOD 1], 2-aminoadenosine was methylated to give 2-amino-2'-O-methyladenosine (Compound 30) and 2-amino-3'-O-methyladenosine (Compound 31).

-Compound 30-

m.p.: 104 - 108 °C (decomposition)

5 NMR(D₂O): δ =3.41(3H,s), 3.81(1H,dd,J=3.4, 12.7Hz), 3.88(1H,dd,J=2.9, 12.7Hz), 4.26(1H,m),
4.49(1H,dd,J=4.9, 6.8Hz), 4.57(1H,dd,J=2.9, 4.9Hz), 5.95(1H,dd,J=6.8Hz), 8.00(1H,s)

-Compound 31-

m.p.: 228 - 228.5 °C

10 NMR(D₂O): δ =3.52(3H,s), 3.80(1H,dd,J=3.4, 12.7Hz), 3.91(1H,dd,J=2.9, 12.7Hz), 4.09(1H,dd,J=2.9,
5.4Hz), 4.35(1H,m), 4.87(1H,dd,J=5.4, 6.4Hz), 5.89(1H,d,J=6.4Hz), 7.99(1H,s)

The following descriptions serve to illustrative pharmaceutical studies of the compounds of the present invention.

(1) Acute toxicity test

15 The test compounds of the present invention, which were dissolved or suspended in 0.5% carboxymethylcellulose (C.M.C.) aqueous solution, were orally administered to groups of 4 to 16 ddY-strain male mice, and the LD₅₀ values were calculated based on the death rate for 7 days thereafter. An example of the results is shown in Table 1.

20 Table 1

25	Test Compound	LD ₅₀ (mg/kg)
30	Compound 1	> 2,000
	Compound 2	> 2,000
35	Compound 3	750 - 1,000
	Compound 5	540
	Compound 9	500 - 1,000
40	Compound 11	800
	Compound 15	500 - 1,000
45	Compound 17	1,000 - 2,000
	Compound 19	1,300
50	Compound 21	about 400
	Compound 22	about 400
55	Compound 24	1,320
	Compound 27	about 400
60	Compound 29	> 2,000

(2) Antihypertensive activity

The test compounds, which were dissolved or suspended in 0.5% C.M.C. aqueous solution, were orally administered to groups of 3 or 4 spontaneously hypertensive rats (SHR), weighing about 300 to 400 g. Blood pressure was measured before and 2,4 and 6 hours after the administration. An example of the results is shown in Table 2. Data show the mean value \pm S.E.

5

10

15

20

25

30

35

40

45

50

55

60

65

Table 2

Test Compound (Dose mg/kg)	Blood pressure (mmHg)			
	Before	2 hrs [Decrease]	4 hrs [Decrease]	6 hrs [Decrease]
Control	219 ± 10	217 ± 12 [1.5 %]	225 ± 17 [-2.6 %]	222 ± 12 [-1.4 %]
Compound 1 (200)	218 ± 21	103 ± 21 [53.2 %]	121 ± 32 [45.8 %]	128 ± 41 [42.8 %]
Compound 2 (200)	227 ± 12	135 ± 28 [41.2 %]	163 ± 40 [29.0 %]	167 ± 47 [27.2 %]
Compound 3 (100)	215 ± 14	102 ± 15 [52.2 %]	106 ± 20 [49.5 %]	111 ± 26 [47.3 %]
Compound 5 (10)	220 ± 11	131 ± 6 [40.2 %]	161 ± 11 [26.5 %]	193 ± 24 [12.7 %]
Compound 9 (30)	221 ± 2	169 ± 7 [23.6 %]	197 ± 5 [10.7 %]	191 ± 9 [13.5 %]
Compound 11 (30)	218 ± 5	189 ± 3 [18.2 %]	206 ± 10 [5.9 %]	204 ± 9 [6.6 %]
Compound 15 (10)	237 ± 8	154 ± 13 [35.2 %]	195 ± 22 [17.7 %]	215 ± 5 [9.1 %]
Compound 17 (30)	231 ± 3	192 ± 15 [17.1 %]	204 ± 5 [11.7 %]	199 ± 8 [14.1 %]
Compound 19 (30)	213 ± 10	122 ± 8 [43.0 %]	155 ± 7 [27.1 %]	157 ± 12 [26.3 %]
Compound 21 (30)	207 ± 6	159 ± 20 [23.9 %]	172 ± 7 [16.9 %]	188 ± 8 [10.8 %]

Compound 22 (10)	227 ± 7	168 ± 6 [25.8 %]	185 ± 7 [18.8 %]	197 ± 3 [13.1 %]	
Compound 24 (10)	226 ± 4	128 ± 7 [43.3 %]	165 ± 7 [27.0 %]	174 ± 4 [23.0 %]	5
Compound 27 (100)	221 ± 9	70 ± 10 [68.4 %]	78 ± 8 [64.8 %]	73 ± 16 [66.9 %]	10
Compound 29 (30)	245 ± 12	131 ± 4 [45.9 %]	195 ± 5 [19.8 %]	204 ± 3 [16.4 %]	15
Adenosine (200)	191 ± 9	202 ± 12 [-6.1 %]	187 ± 3 [1.8 %]	198 ± 4 [-3.2 %]	20

As shown by the above-mentioned results, the compounds of the present invention have excellent hypotensive effects even at low doses. However, adenosine does not decrease blood pressure at oral administration.

Bradycardia was observed at the administration of adenosine or 2-substituted adenosines, however, the compounds of the present invention O-alkylated at the 2-or 3-position of ribose did not show such a side effect.

Thus the compounds of the present invention have excellent antihypertensive effect at oral administration and do not have side effect such as bradycardia. Therefore, the compounds of the present invention are not only useful as antihypertensive drugs for oral administration, but also as drugs for various diseases caused by hypertension, e.g., cerebrovascular disease such as cerebral hemorrhage, cerebral infarction or subarachnoidal hemorrhage, cardiopathy such as congestive heart failure, myocardial infarction or sudden cardiac death, and renal insufficiency.

The compounds of the present invention can be made into pharmaceutical compositions by combination with appropriate medicinal carriers or diluents, and can be formulated into preparations in solid, semisolid, liquid or gaseous form such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, aerosols and cataplasms in usual ways for oral or parenteral administrations.

In pharmaceutical dosage forms, the compounds of the present invention can be used in the form of their pharmaceutically acceptable salts, and also can be used alone or in appropriate association, as well as in combination with other pharmaceutically active components.

In case of oral preparations, the compounds can be used alone or combined with appropriate additives to make tablets, powders, granules or capsules, e.g. with conventional additives such as lactose, mannitol, corn starch or potato starch; with binders such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins; with disintegrators such as corn starch, potato starch or sodium carboxymethylcellulose; with lubricants such as talc or magnesium stearate; and if desired, with diluents, buffering agents, moistening agents, preservatives and flavoring agents.

The compounds of the invention can also be made into an ointment by combination with an ointment base such as vaseline, paraffin, plastibase, simple ointment, hydrophilic ointment, hydrophilic vaseline or hydrophilic plastibase.

Furthermore, the compounds of the invention can be made into a suppository by mixing with a variety of bases, e.g. fatty and oily base such as cacao butter, emulsifying base or water-soluble base such as macrogol.

The compounds of the present invention can be formulated into a preparations for injections by dissolving, suspending or emulsifying in aqueous or non-aqueous solvent, such as distilled water for injection, physiologically saline solution, vegetable oil, synthetic aliphatic acid glycerides, esters of higher aliphatic acid or propylene glycol.

In case of inhalations or aerosol preparations, the compounds of the invention in the form of a liquid or minute powder can be filled up in an aerosol container with gas or liquid spraying agent, and if desired, with conventional adjuvants such as humidifying agents or dispersing agent. They can also be used as pharmaceuticals for a non-pressurized preparation such as in a nebulizer or an atomizer.

Cataplasms can be prepared by mixing the compounds with mentha oil, concentrated glycerin, kaolin or

other suitable additives.

The desirable dose of the compounds of the present invention varies with the subject, form of the drug, method and period of administration. However, in order to obtain desirable effects, generally it is recommended to administer orally 0.2 to 5,000 mg, preferably 1 to 3,000 mg daily. Unit preparations are also recommended for administration in one to several units daily.

In case of parenteral administrations e.g. injections, doses of the compounds in the order of one tenth to one second of the above dose are preferable as daily doses.

Some prescriptions of the pharmaceutical compositions are shown below as examples which contain the compounds of the present invention as active ingredients.

Prescription example 1 (tablet)

Component	Content in a tablet (mg)
compound of this invention	25
lactose	175
corn starch	40
magnesium stearate	10
Total 250 mg	

Prescription example 2 (capsule)

Component	Content in a capsule (mg)
compound of this invention	50
lactose	250
Total 300 mg	

Prescription example 3 (inhalation)

Component	Content in a inhalation (g)
compound of this invention	1
lactose	5
<hr/>	
Total	6 g

Prescription example 4 (injection)


Component	Content in an ampule (mg)
compound of this invention	10
sodium chloride	proper amount
distilled water for injection	proper amount
<hr/>	
Total	1 ml

Prescription example 5 (ointment)

Component	Weight (g)
compound of this invention	1
emulsified wax	30
white petrolatum	50
liquid paraffin	20
<hr/>	
Total	101 g

Claims

1. An adenosine derivative having the formula (I):



wherein each of R_1 , R_2 and R_3 , which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl group;

or pharmaceutically acceptable salt thereof.

2. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 1 wherein R_1 is a lower alkyl group.

3. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 2 wherein X is a lower alkyl group.

4. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 2 wherein Y is a lower alkyl group.


5. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 1 wherein R_3 is a lower alkyl group.

6. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 5 wherein X is a lower alkyl group.

7. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 5 wherein Y is a lower alkyl group.

8. An adenosine derivative or pharmaceutically acceptable salt thereof according to claim 1 wherein X is halogen.

9. An antihypertensive composition comprising as an active ingredient an effective amount of at least one adenosine derivative of the formula (I):




(I)

wherein each of R_1 , R_2 and R_3 , which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl group;
or pharmaceutically acceptable salt thereof.


10. An antihypertensive composition according to claim 9, which is formulated into a form suitable for oral administration.

11. A process for preparing an adenosine derivative of the formula (I):



(I)

wherein each of R_1 , R_2 and R_3 , which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl group;
which comprises O-alkylating a compound of the formula (II):



(II)

wherein each of R₁', R₂' and R₃', which may be the same or different, is hydrogen, a lower alkyl group or a protecting group; X' is hydrogen, a lower alkyl group, an amino group or halogen; and Y' is an amino group, a lower alkylamino group or halogen; and then, if desired, carrying out an amination or an alkylation of the 6-position and a removal of a protecting group.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 269 574
A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **87810687.1**(51) Int. Cl.4: **C07H 19/173 , A61K 31/70**(22) Date of filing: **23.11.87**(30) Priority: **27.11.86 JP 283904/86**(43) Date of publication of application:
01.06.88 Bulletin 88/22(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE(88) Date of deferred publication of the search report:
22.03.89 Bulletin 89/12(71) Applicant: **NIPPON ZOKI PHARMACEUTICAL CO. LTD.**

**10, 2-chome Hirano-machi Higashi-ku
Osaka-shi Osaka-fu(JP)**


(72) Inventor: **Yamada, Toshio c/o Institute of Bio-Active Science**

**Nippon Zoki Pharm. Co. Ltd. 442-1, Kinashi
Yashiro-cho Katoh-gun Hyogo(JP)**

Inventor: **Kageyama, Ken-ichi Institute of Bio-Active Science**
**Nippon Zoki Pharm. Co. Ltd. 442-1, Kinashi
Yashiro-cho Katoh-gun Hyogo(JP)**

(74) Representative: **Kerr, Andrew**
Postfach 122 Finkelerweg 44
CH-4144 Arlesheim BL(CH)(54) **Novel adenosine derivatives and pharmaceutical compositions containing them as an active ingredient.**

(57) The present invention relates to novel adenosine derivatives having the formula (I):



(I)

wherein each of R_1 , R_2 and R_3 , which may be the same or different, is hydrogen or a lower alkyl group; X is hydrogen, a lower alkyl group, an amino group or halogen; and Y is hydrogen or a lower alkyl

EP 0 269 574 A3

group;

and pharmaceutically acceptable salt thereof. These compounds are useful as antihypertensive agents, for example, for the treatments of hypertension and various diseases caused by hypertension, such as cerebrovascular diseases, cardiopathy or renal insufficiency.



EP 87 81 0687

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	GB-A-1 367 785 (TAKEDA YAKUHHIN KOGYO K.K.) * Examples 6,7; page 1, lines 15-21 * ---	1,3,9	C 07 H 19/173 A 61 K 31/70
X	FR-A-1 324 011 (FARBWERKE HOECHST AG) * Example 1; page 3, left-hand column, last paragraph * ---	1,9	
X	GB-A-1 240 087 (BOEHRINGER MANNHEIM GmbH) * Whole document * ---	1,4,9,10	
X	CHEMICAL ABSTRACTS, vol. 92, 1980, page 138, no. 105074e, Columbus, Ohio, US; N.J. CUSACK et al.: "Effects of D- and L-enantiomers of adenosine, AMP and ADP and their 2-chloro- and 2-azido-analogs on human platelets", & PROC. R. SOC. LONDON, SER. B 1979, 206(1163), 139-44 * Abstract * ---	1,8	
X	GB-A-1 139 160 (C.F. BOEHRINGER & SOEHNE GmbH) * Formula II, "starting products"; examples 1,2,10 * ---	1,4	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 07 H 19/00
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 15, no. 4, 1972, pages 387-390, Columbus, Ohio, US; K. KIKUGAWA et al.: "Platelet aggregation inhibitors. 2. Inhibition of platelet aggregation by 5'-,2-,6-, and 8-substituted adenosines" * Page 387, compounds 5c,5g * --- -/-	1,4,9	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12-12-1988	Examiner VAN BIJLEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 13, no. 3, 1970, pages 427-430, Columbus, Ohio, US; J.A. MONTGOMERY et al.: "2-Fluoropurine ribonucleosides" * Page 427, "(2-fluoradenosine), compound 11 * ---	1,4,8,9	
X	CHEMICAL ABSTRACTS, vol. 72, 1970, page 352, no. 21921c, Columbus, Ohio, US; & SA-A-68 05 477 (BOEHRINGER, C.F., UND SOEHNE GmbH) 28-01-1969 * Abstract * ---	1,4	
X	CHEMICAL ABSTRACTS, vol. 84, 1976, page 577, no. 59957e, Columbus, Ohio, US; & JP-A-75 121 297 (KOHJIN CO., LTD) 23-09-1975 * Abstract * ---	1,8	
X	CHEMICAL ABSTRACTS, vol. 97, 1982, page 23, no. 120125u, Columbus, Ohio, US; M.L. PAUL et al.: "The influence of glycosidic conformation and charge distribution on activity of adenine nucleosides as presynaptic inhibitors of acetylcholine release", & J. PHARMACOL. EXP. THER. 1982, 222(1), 241-5 * Abstract * --- -/-	1,8,9	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 12-12-1988	Examiner VAN BIJLEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	CHEMICAL ABSTRACTS, vol. 97, 1982, page 751, no. 24157k, Columbus, Ohio, US; M.J. ROBINS et al.: "Nucleic acid related compounds. 36. Synthesis of the 2'-O-methyl and 3'-O-methyl ethers of guanosine and 2-aminoadenosine and correlation of O'-methylnucleoside carbon-13 NMR spectral shifts", & CAN. J. CHEM. 1981, 59(24), 3360-4 * Abstract * ---	1,5	
X	CHEMICAL ABSTRACTS, vol. 103, 1985, page 708, no. 196359m, Columbus, Ohio, US; A. MATSUDA et al.: "Palladium-catalyzed cross-coupling of 2-iodoadenosine with terminal alkynes: synthesis and biological activities of 2-alkynyladenosines", & CHEM. PHARM. BULL. 1985, 33(4), 1766-9 * Abstract * ---	1,8	
X	BIOCHEMISTRY, vol. 15, no. 13, 1976, pages 2735-2740, Columbus, Ohio, US; Z. KAZIMIERCZUK et al.: "Methylation of adenosine in strongly alkaline medium: preparation and properties of O'-methyl; Derivatives of adenosine and N6-methyladenosine" * Pages 2337-2339 * ---	1,2,4,5	
X	BIOCHEMISTRY, vol. 15, no. 23, 1976, pages 5052-5057; B. SINGER et al.: "Alkylation of ribose in RNA reacted with ethylnitrosourea at neutrality" * Page 5053 * --- -/-	1,5	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12-12-1988	Examiner VAN BIJLEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	JOURNAL OF ORGANIC CHEMISTRY, vol. 39, no. 25, 1974, pages 3674-3676, Columbus, Ohio, US; L.M. PIKE et al.: "Mixed alkylation (Methylation and ethylation) of adenosine by diazoethane in aqueous 1,2-dimethoxyethane" * Whole article *	1,4,5	
X	JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 89, no. 14, 5th July 1967, pages 3612-3622, Columbus, Ohio, US; A.D. BROOM et al.: "Interaction and association of bases and nucleosides in aqueous solutions. V. Studies of the association of purine nucleosides by vapor pressure osmometry and by proton magnetic resonance" * Page 3613, table I *	1,4,5	
X	CHEMICAL ABSTRACTS, vol. 84, 1976, page 135, no. 1373v, Columbus, Ohio, US; C.-M. WEI et al.: "N ⁶ ,O ^{2'} -dimethyladenosine, a novel methylated ribonucleoside next to the 5'-terminal of animal cell and virus mRNAs", & NATURE (LONDON) 1975, 257(5523), 251-3 * Abstract *	1,4,5	
X	BIOCHEMISTRY, vol. 7, no. 4, April 1968, pages 1413-1420, Columbus, Ohio, US; J.B. GIN et al.: "The preparation and properties of O-methylated adenosine derivatives" * Page 1417, table I *	1,5	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 12-12-1988	Examiner VAN BIJLEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
E	EP-A-0 253 962 (HOECHST-JAPAN LTD) * Pages 5,6; claims * -----		
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12-12-1988	Examiner VAN BIJLEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			